

# Representations of the restricted enveloping algebra $u(\mathfrak{m})$ in characteristic 2

Dirceu Bagio

dirceu.bagio@ufsm.br

Federal University of Santa Maria

Cimpa School, Florianópolis

February 7, 2021

\* This is a work in progress with N. Andruskiewitsch, S. D. Flora and D. Flores.

## Outline

- ▶ The Drinfeld double  $D(H)$  of the restricted Jordan plane.

## Outline

- ▶ The Drinfeld double  $D(H)$  of the restricted Jordan plane.
- ▶  $D(H)$  as an abelian extension:

## Outline

- ▷ The Drinfeld double  $D(H)$  of the restricted Jordan plane.
- ▷  $D(H)$  as an abelian extension:

$$\mathbf{T} \xhookrightarrow{\iota} D(H) \twoheadrightarrow^{\pi} u(\mathfrak{m})$$

- ▷  $u(\mathfrak{m})$  is of tame representation type.

## Outline

- ▷ The Drinfeld double  $D(H)$  of the restricted Jordan plane.
- ▷  $D(H)$  as an abelian extension:

$$\mathbf{T} \xhookrightarrow{\iota} D(H) \twoheadrightarrow^{\pi} u(\mathfrak{m})$$

- ▷  $u(\mathfrak{m})$  is of tame representation type.
- ▷ Irreducible representations of  $u(\mathfrak{m})$ .

In what follows,  $\mathbb{k}$  is an algebraically closed field of characteristic 2.

In what follows,  $\mathbb{k}$  is an algebraically closed field of characteristic 2.

## Restricted Jordan plane

It is the algebra  $\mathcal{B}$  of dimension  $2^4$  presented by generators  $x_1, x_2$  with defining relations

$$x_1^2 = 0, \quad x_2^2 x_1 = x_1 x_2^2 + x_1 x_2 x_1, \quad (1)$$

$$x_2^4 = 0, \quad x_1 x_2 x_1 x_2 = x_2 x_1 x_2 x_1. \quad (2)$$

In what follows,  $\mathbb{k}$  is an algebraically closed field of characteristic 2.

## Restricted Jordan plane

It is the algebra  $\mathcal{B}$  of dimension  $2^4$  presented by generators  $x_1, x_2$  with defining relations

$$x_1^2 = 0, \quad x_2^2 x_1 = x_1 x_2^2 + x_1 x_2 x_1, \quad (1)$$

$$x_2^4 = 0, \quad x_1 x_2 x_1 x_2 = x_2 x_1 x_2 x_1. \quad (2)$$

## Bosonization

Let  $\Gamma = \langle g \rangle$  be the cyclic group of order 2, written multiplicatively. The bosonization  $H := \mathcal{B} \# \mathbb{k}\Gamma$  is a pointed Hopf algebra of dimension  $2^5$  generated by  $x_1, x_2, g$  with satisfies the previous relations and

$$gx_1 = x_1g, \quad gx_2 = x_2g + x_1g, \quad g^2 = 1. \quad (3)$$

The coproduct of  $H$  is given by

$$\Delta(g) = g \otimes g, \quad \Delta(x_i) = x_i \otimes 1 + g \otimes x_i, \quad i \in \mathbb{I}_2.$$



## Drinfeld double of $H$

The Drinfeld double of  $H$  is  $D(H) = H \otimes H^{*\text{op}}$  as coalgebra. As algebra,  $D(H)$  is generated by  $x_1, x_2, g, w_1, w_2, \gamma$  with relations (1), (2), (3) and

## Drinfeld double of $H$

The Drinfeld double of  $H$  is  $D(H) = H \otimes H^{*\text{op}}$  as coalgebra. As algebra,  $D(H)$  is generated by  $x_1, x_2, g, w_1, w_2, \gamma$  with relations (1), (2), (3) and

$$w_1^2 = 0,$$

$$w_2^2 w_1 = w_1 w_2^2 + w_1 w_2 w_1,$$

$$w_2^4 = 0,$$

$$w_1 w_2 w_1 w_2 = w_2 w_1 w_2 w_1,$$

$$\gamma^2 = \gamma,$$

$$w_i \gamma = \gamma w_i + w_i,$$

$$w_1 x_1 = x_1 w_1,$$

$$w_1 x_2 = x_2 w_1 + 1 + g,$$

$$w_1 g = g w_1,$$

$$w_2 x_1 = x_1 (w_1 + w_2) + 1 + g,$$

$$w_2 g = g (w_1 + w_2),$$

$$\gamma x_i = x_i \gamma + x_i,$$

$$w_2 x_2 = x_2 w + g \gamma,$$

## Drinfeld double of $H$

The Drinfeld double of  $H$  is  $D(H) = H \otimes H^{*\text{op}}$  as coalgebra. As algebra,  $D(H)$  is generated by  $x_1, x_2, g, w_1, w_2, \gamma$  with relations (1), (2), (3) and

$$w_1^2 = 0,$$

$$w_2^2 w_1 = w_1 w_2^2 + w_1 w_2 w_1,$$

$$w_2^4 = 0,$$

$$w_1 w_2 w_1 w_2 = w_2 w_1 w_2 w_1,$$

$$\gamma^2 = \gamma,$$

$$w_i \gamma = \gamma w_i + w_i,$$

$$w_1 x_1 = x_1 w_1,$$

$$w_1 x_2 = x_2 w_1 + 1 + g,$$

$$w_1 g = g w_1,$$

$$w_2 x_1 = x_1 (w_1 + w_2) + 1 + g,$$

$$w_2 g = g (w_1 + w_2), \quad \gamma x_i = x_i \gamma + x_i,$$

$$w_2 x_2 = x_2 w + g \gamma,$$

We have  $\dim D(H) = 2^{10}$ .

Fix the following elements in  $D(H)$ :

$$x_{21} = x_1 x_2 + x_2 x_1, \quad w_{21} = w_1 w_2 + w_2 w_1.$$

Fix the following elements in  $D(H)$ :

$$x_{21} = x_1 x_2 + x_2 x_1, \quad w_{21} = w_1 w_2 + w_2 w_1.$$

## Central Hopf subalgebra

The subalgebra  $\mathbf{T}$  of  $D(H)$  generated by  $x_1$ ,  $x_{21}$ ,  $w_1$ ,  $w_{21}$  and  $g$  is a normal local commutative Hopf subalgebra with defining relations

$$x_1^2 = 0, \quad x_{21}^2 = 0, \quad w_1^2 = 0, \quad w_{21}^2 = 0, \quad g^2 = 1.$$

Also  $\dim \mathbf{T} = 2^5$ .

Fix the following elements in  $D(H)$ :

$$x_{21} = x_1 x_2 + x_2 x_1, \quad w_{21} = w_1 w_2 + w_2 w_1.$$

## Central Hopf subalgebra

The subalgebra  $\mathbf{T}$  of  $D(H)$  generated by  $x_1$ ,  $x_{21}$ ,  $w_1$ ,  $w_{21}$  and  $g$  is a normal local commutative Hopf subalgebra with defining relations

$$x_1^2 = 0, \quad x_{21}^2 = 0, \quad w_1^2 = 0, \quad w_{21}^2 = 0, \quad g^2 = 1.$$

Also  $\dim \mathbf{T} = 2^5$ .

Hence

$$\mathbf{T} \xhookrightarrow{\iota} D(H) \twoheadrightarrow^{\pi} D(H)/D(H)\mathbf{T}^+$$

is an exact sequence of Hopf algebras.

We fix the following elements of  $D(H)/D(H)\mathbf{T}^+$ :

$$a = \bar{x}_2, \quad b = \bar{w}_2, \quad c = \bar{\gamma}.$$

We fix the following elements of  $D(H)/D(H)\mathbf{T}^+$ :

$$a = \bar{x}_2, \quad b = \bar{w}_2, \quad c = \bar{\gamma}.$$

## Hopf algebra quotient

The algebra  $D(H)/D(H)\mathbf{T}^+$  is generated by  $a, b, c$  and satisfies the relations

$$\begin{aligned} ab + ba &= c, & ac + ca &= a, & bc + cb &= b, \\ a^4 &= b^4 = 0, & c^2 + c &= 0. \end{aligned}$$



We fix the following elements of  $D(H)/D(H)\mathbf{T}^+$ :

$$a = \bar{x}_2, \quad b = \bar{w}_2, \quad c = \bar{\gamma}.$$

## Hopf algebra quotient

The algebra  $D(H)/D(H)\mathbf{T}^+$  is generated by  $a, b, c$  and satisfies the relations

$$\begin{aligned} ab + ba &= c, & ac + ca &= a, & bc + cb &= b, \\ a^4 &= b^4 = 0, & c^2 + c &= 0. \end{aligned}$$

The Hopf algebra  $D(H)/D(H)\mathbf{T}^+$  is a well-known algebra in modular Lie theory.

Denote by  $\mathfrak{s}$  the unique, up to isomorphism, simple Lie algebra of dimension 3, that is,  $\mathfrak{s}$  has a basis  $\{e, f, h\}$  and bracket

$$[e, f] = h, \quad [e, h] = e, \quad [f, h] = f.$$

Denote by  $\mathfrak{s}$  the unique, up to isomorphism, simple Lie algebra of dimension 3, that is,  $\mathfrak{s}$  has a basis  $\{e, f, h\}$  and bracket

$$[e, f] = h, \quad [e, h] = e, \quad [f, h] = f.$$

The Lie algebra  $\mathfrak{s}$  is not restricted. The minimal 2-envelope of  $\mathfrak{s}$  is a 5-dimensional Lie algebra  $\mathfrak{m}$ .

Denote by  $\mathfrak{s}$  the unique, up to isomorphism, simple Lie algebra of dimension 3, that is,  $\mathfrak{s}$  has a basis  $\{e, f, h\}$  and bracket

$$[e, f] = h, \quad [e, h] = e, \quad [f, h] = f.$$

The Lie algebra  $\mathfrak{s}$  is not restricted. The minimal 2-envelope of  $\mathfrak{s}$  is a 5-dimensional Lie algebra  $\mathfrak{m}$ .

### The algebra $u(\mathfrak{m})$

The restricted enveloping algebra  $u(\mathfrak{m})$  of  $\mathfrak{m}$  is isomorphic to  $D(H)/D(H)\mathbf{T}^+$  via

$$e \mapsto a, \quad f \mapsto b, \quad h \mapsto c.$$

and we have an exact sequence of Hopf algebras

$$\mathbf{T} \xhookrightarrow{\iota} D(H) \twoheadrightarrow^{\pi} u(\mathfrak{m})$$

Denote by  $\mathfrak{s}$  the unique, up to isomorphism, simple Lie algebra of dimension 3, that is,  $\mathfrak{s}$  has a basis  $\{e, f, h\}$  and bracket

$$[e, f] = h, \quad [e, h] = e, \quad [f, h] = f.$$

The Lie algebra  $\mathfrak{s}$  is not restricted. The minimal 2-envelope of  $\mathfrak{s}$  is a 5-dimensional Lie algebra  $\mathfrak{m}$ .

### The algebra $u(\mathfrak{m})$

The restricted enveloping algebra  $u(\mathfrak{m})$  of  $\mathfrak{m}$  is isomorphic to  $D(H)/D(H)\mathbf{T}^+$  via

$$e \mapsto a, \quad f \mapsto b, \quad h \mapsto c.$$

and we have an exact sequence of Hopf algebras

$$\mathbf{T} \xhookrightarrow{\iota} D(H) \twoheadrightarrow^{\pi} u(\mathfrak{m})$$

For this reason we are interested in the representations of  $u(\mathfrak{m})$ .

Let  $V_0$ , respectively  $V_1$ , denote the one-dimensional  $\mathfrak{u}(\mathfrak{m})$ -module, respectively the three dimensional  $\mathfrak{u}(\mathfrak{m})$ -module  $\mathfrak{s}$  with the adjoint representation  $\text{ad}$ .

Let  $V_0$ , respectively  $V_1$ , denote the one-dimensional  $\mathfrak{u}(\mathfrak{m})$ -module, respectively the three dimensional  $\mathfrak{u}(\mathfrak{m})$ -module  $\mathfrak{s}$  with the adjoint representation  $\text{ad}$ .


Thus  $V_1$  in the basis  $\{v_1, v_2, v_3\} := \{b, c, a\}$  of  $\mathfrak{s}$  is given by  $\text{ad } a = A$ ,  $\text{ad } b = B$ ,  $\text{ad } c = C$ , where

$$A = \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}, \quad B = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}, \quad C = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

### Theorem.

The simple modules of  $\mathfrak{u}(\mathfrak{m})$  are  $V_0$  and  $V_1$ .


Consider the following 8-dimensional  $u(\mathfrak{m})$ -module  $M$



where the arrows oriented from left to right indicate the action of  $a$  while the arrows from right to left are the action of  $b$ .



Consider the following 8-dimensional  $\mathfrak{u}(\mathfrak{m})$ -module  $M$




where the arrows oriented from left to right indicate the action of  $a$  while the arrows from right to left are the action of  $b$ .


**Theorem.**

- ↪ The projective cover of the simple module  $V_0$  is  $P(V_0) = M$ .
- ↪  $P(V_0) \simeq \mathfrak{u}(\mathfrak{m})e_0$ , where  $e_0 = (1 + ab + a^2b^2)(1 + c)$  is a primitive idempotent of  $\mathfrak{u}(\mathfrak{m})$ .

Consider the following 8-dimensional  $\mathfrak{u}(\mathfrak{m})$ -module  $N$



Consider the following 8-dimensional  $u(\mathfrak{m})$ -module  $N$



**Theorem.**

- ↪ The projective cover of the simple module  $V_1$  is  $P(V_1) = N$ .
- ↪  $P(V_1) \simeq u(\mathfrak{m})e_1$ , where  $e_1 = (1 + a^2b^2)c$  is a primitive idempotent of  $u(\mathfrak{m})$ .

Let  $e = e_0 + e_1$  where  $e_0$  and  $e_1$  are the primitive idempotents in  $\mathfrak{u}(\mathfrak{m})$  generating the projective covers  $P(V_0)$  and  $P(V_1)$ . Then the basic algebra associated to  $\mathfrak{u}(\mathfrak{m})$  is

$$\mathfrak{u}(\mathfrak{m})^b = e\mathfrak{u}(\mathfrak{m})e.$$

Let  $e = e_0 + e_1$  where  $e_0$  and  $e_1$  are the primitive idempotents in  $\mathfrak{u}(\mathfrak{m})$  generating the projective covers  $P(V_0)$  and  $P(V_1)$ . Then the basic algebra associated to  $\mathfrak{u}(\mathfrak{m})$  is

$$\mathfrak{u}(\mathfrak{m})^b = e\mathfrak{u}(\mathfrak{m})e.$$

The basic algebra  $\mathfrak{u}(\mathfrak{m})^b$  has a basis

$$\{e_0, e_1, ae_0, a^3e_1, b^3e_0, be_1, a^3b^3e_0, abe_1\}.$$


Let  $e = e_0 + e_1$  where  $e_0$  and  $e_1$  are the primitive idempotents in  $u(\mathfrak{m})$  generating the projective covers  $P(V_0)$  and  $P(V_1)$ . Then the basic algebra associated to  $u(\mathfrak{m})$  is

$$u(\mathfrak{m})^b = eu(\mathfrak{m})e.$$

The basic algebra  $u(\mathfrak{m})^b$  has a basis

$$\{e_0, e_1, ae_0, a^3e_1, b^3e_0, be_1, a^3b^3e_0, abe_1\}.$$

The ordinary quiver of  $\mathfrak{u}(\mathfrak{m})^b$ , denoted by  $Q := Q_{\mathfrak{u}(\mathfrak{m})^b}$ , is



## Theorem.

We have that  $\mathfrak{u}(\mathfrak{m})^b \simeq \mathbb{K}Q/I$ , where

$$I = \langle \alpha_1\beta_1, \alpha_2\beta_2, \beta_1\alpha_1, \beta_2\alpha_2, \alpha_1\beta_2 + \alpha_2\beta_1, \beta_1\alpha_2 + \beta_2\alpha_1 \rangle$$

is the kernel of the algebra epimorphism  $\varphi : \mathbb{K}Q \rightarrow \mathfrak{u}(\mathfrak{m})^b$  defined by

$$\varphi(\alpha_1) = a^3 e_1, \quad \varphi(\alpha_2) = b e_1, \quad \varphi(\beta_1) = a e_0, \quad \varphi(\beta_2) = b^3 e_0,$$

and  $\varphi(\varepsilon_i) = e_i$ .

## Theorem.

We have that  $\mathfrak{u}(\mathfrak{m})^b \simeq \mathbb{k}Q/I$ , where

$$I = \langle \alpha_1\beta_1, \alpha_2\beta_2, \beta_1\alpha_1, \beta_2\alpha_2, \alpha_1\beta_2 + \alpha_2\beta_1, \beta_1\alpha_2 + \beta_2\alpha_1 \rangle$$

is the kernel of the algebra epimorphism  $\varphi : \mathbb{k}Q \rightarrow \mathfrak{u}(\mathfrak{m})^b$  defined by

$$\varphi(\alpha_1) = a^3 e_1, \quad \varphi(\alpha_2) = b e_1, \quad \varphi(\beta_1) = a e_0, \quad \varphi(\beta_2) = b^3 e_0,$$

and  $\varphi(\varepsilon_i) = e_i$ .

## Corollary

$\mathfrak{u}(\mathfrak{m})^b$  is a special biserial algebra. Particularly,  $\mathfrak{u}(\mathfrak{m})^b$  is tame representation type.



## Theorem.

We have that  $u(\mathfrak{m})^b \simeq \mathbb{k}Q/I$ , where

$$I = \langle \alpha_1\beta_1, \alpha_2\beta_2, \beta_1\alpha_1, \beta_2\alpha_2, \alpha_1\beta_2 + \alpha_2\beta_1, \beta_1\alpha_2 + \beta_2\alpha_1 \rangle$$

is the kernel of the algebra epimorphism  $\varphi : \mathbb{k}Q \rightarrow u(\mathfrak{m})^b$  defined by

$$\varphi(\alpha_1) = a^3 e_1, \quad \varphi(\alpha_2) = b e_1, \quad \varphi(\beta_1) = a e_0, \quad \varphi(\beta_2) = b^3 e_0,$$

and  $\varphi(\varepsilon_i) = e_i$ .

## Corollary

$u(\mathfrak{m})^b$  is a special biserial algebra. Particularly,  $u(\mathfrak{m})^b$  is tame representation type.

## Corollary

Since  $u(\mathfrak{m})$  is Morita equivalent to  $u(\mathfrak{m})^b$  it follows that  $u(\mathfrak{m})$  is tame representation type.

→ The classification of all indecomposable modules of a special biserial algebra was given in [4, Proposition 2.3].

- ~> The classification of all indecomposable modules of a special biserial algebra was given in [4, Proposition 2.3].
- ~> They are either string modules or band modules.

- ~> The classification of all indecomposable modules of a special biserial algebra was given in [4, Proposition 2.3].
- ~> They are either string modules or band modules.
- ~> Determining strings and bands in  $\mathbb{k}Q/I$  is a combinatorial problem.

- ~> The classification of all indecomposable modules of a special biserial algebra was given in [4, Proposition 2.3].
- ~> They are either string modules or band modules.
- ~> Determining strings and bands in  $\mathbb{k}Q/I$  is a combinatorial problem.

In our case, for instance, consider words in the vocabulary  $\{\alpha_i^{\pm 1}, \beta_i^{\pm 1} : i = 1, 2\}$ .

- ~> The classification of all indecomposable modules of a special biserial algebra was given in [4, Proposition 2.3].
- ~> They are either string modules or band modules.
- ~> Determining strings and bands in  $\mathbb{k}Q/I$  is a combinatorial problem.

In our case, for instance, consider words in the vocabulary  $\{\alpha_i^{\pm 1}, \beta_i^{\pm 1} : i = 1, 2\}$ .

The words  $a = \alpha_1 \alpha_1^{-1} \alpha_2$  and  $b = \alpha_1 \beta_2$  are not string. In fact,  $a$  is not a string because  $\alpha_1 \alpha_1^{-1}$  is a "piece" of  $a$  and  $b$  is not a string because  $\alpha_1 \beta_2$  is a monomial of the binomial relation  $\alpha_1 \beta_2 + \alpha_2 \beta_1$ .

- ~> The classification of all indecomposable modules of a special biserial algebra was given in [4, Proposition 2.3].
- ~> They are either string modules or band modules.
- ~> Determining strings and bands in  $\mathbb{k}Q/I$  is a combinatorial problem.

In our case, for instance, consider words in the vocabulary  $\{\alpha_i^{\pm 1}, \beta_i^{\pm 1} : i = 1, 2\}$ .

The words  $a = \alpha_1 \alpha_1^{-1} \alpha_2$  and  $b = \alpha_1 \beta_2$  are not string. In fact,  $a$  is not a string because  $\alpha_1 \alpha_1^{-1}$  is a "piece" of  $a$  and  $b$  is not a string because  $\alpha_1 \beta_2$  is a monomial of the binomial relation  $\alpha_1 \beta_2 + \alpha_2 \beta_1$ .

The words  $s = \alpha_1 \alpha_2^{-1} \alpha_1$  and  $t = \beta_1^{-1} \beta_2$  are examples of strings.

## String

Consider the words  $s_1 = \alpha_1 \alpha_2^{-1}$ ,  $s_2 = \alpha_1^{-1} \alpha_2$ ,  $s_3 = \beta_1 \beta_2^{-1}$  and  $s_4 = \beta_1^{-1} \beta_2$  and  $r$  an integer. The families of string in  $Q$  are the following:

$$w_1(r) = s_1^r, \quad w_2(r) = s_2^r, \quad r \geq 1,$$

$$w_3(r) = s_3^r, \quad w_4(r) = s_4^r, \quad r \geq 1,$$

$$w_5(r) = s_1^r \alpha_1, \quad w_6(r) = (s_1^{-1})^r \alpha_2, \quad r \geq 0$$

$$w_7(r) = s_3^r \beta_1, \quad w_8(r) = (s_3^{-1})^r \beta_2, \quad r \geq 0.$$



## String

Consider the words  $s_1 = \alpha_1 \alpha_2^{-1}$ ,  $s_2 = \alpha_1^{-1} \alpha_2$ ,  $s_3 = \beta_1 \beta_2^{-1}$  and  $s_4 = \beta_1^{-1} \beta_2$  and  $r$  an integer. The families of string in  $Q$  are the following:

$$\begin{aligned} w_1(r) &= s_1^r, & w_2(r) &= s_2^r, & r &\geq 1, \\ w_3(r) &= s_3^r, & w_4(r) &= s_4^r, & r &\geq 1, \\ w_5(r) &= s_1^r \alpha_1, & w_6(r) &= (s_1^{-1})^r \alpha_2, & r &\geq 0, \\ w_7(r) &= s_3^r \beta_1, & w_8(r) &= (s_3^{-1})^r \beta_2, & r &\geq 0. \end{aligned}$$

Similarly, we have the notion of band. For our case, there are 2 families of band in  $Q$ .

In order to illustrate how to associate an indecomposable module to a string, we consider the string  $w_1(1) = s_1 = \alpha_1 \alpha_2^{-1}$ :

$$1 \xrightarrow{\alpha_1} 2 \xleftarrow{\alpha_2} 1$$

In order to illustrate how to associate an indecomposable module to a string, we consider the string  $w_1(1) = s_1 = \alpha_1 \alpha_2^{-1}$ :

$$1 \xrightarrow{\alpha_1} 2 \xleftarrow{\alpha_2} 1$$

The right  $\mathbb{k}Q/I$ -module  $U(w_1(1)) := \mathbb{k}\{w_1, w_2, w_3\}$  (a vector for each vertex) is given by:

$$u_1 \cdot \epsilon_1 = u_1, \quad u_1 \cdot \epsilon_2 = 0, \quad u_1 \cdot \alpha_1 = u_2, \quad u_1 \cdot \alpha_2 = 0,$$

$$u_2 \cdot \epsilon_1 = 0, \quad u_2 \cdot \epsilon_2 = u_2, \quad u_2 \cdot \alpha_1 = 0, \quad u_2 \cdot \alpha_2 = 0,$$

$$u_3 \cdot \epsilon_1 = u_3, \quad u_3 \cdot \epsilon_2 = 0, \quad u_3 \cdot \alpha_1 = 0, \quad u_3 \cdot \alpha_2 = u_2,$$

The algebra isomorphism  $u(\mathfrak{m})^b \simeq \mathbb{k}Q/I$  and an anti-isomorphism of Hopf algebras  $u(\mathfrak{m})^b \rightarrow u(\mathfrak{m})^b$  induce on  $U(w_1(1))$  a left  $u(\mathfrak{m})^b$ -module structure via

The algebra isomorphism  $u(\mathfrak{m})^b \simeq \mathbb{k}Q/I$  and an anti-isomorphism of Hopf algebras  $u(\mathfrak{m})^b \rightarrow u(\mathfrak{m})^b$  induce on  $U(w_1(1))$  a left  $u(\mathfrak{m})^b$ -module structure via

$$e_1 \cdot u_1 = u_1, \quad e_2 \cdot u_1 = 0, \quad ae_1 \cdot u_1 = u_2, \quad b^3 e_1 \cdot eu_1 = 0,$$

$$e_1 \cdot u_2 = 0, \quad e_2 \cdot u_2 = u_2, \quad ae_1 \cdot u_2 = 0, \quad b^3 e_1 \cdot u_2 = 0,$$

$$e_1 \cdot u_3 = u_3, \quad e_2 \cdot u_3 = 0, \quad ae_1 \cdot u_3 = 0, \quad b^3 e_1 \cdot u_3 = u_2.$$

The algebra isomorphism  $u(\mathfrak{m})^b \simeq \mathbb{k}Q/I$  and an anti-isomorphism of Hopf algebras  $u(\mathfrak{m})^b \rightarrow u(\mathfrak{m})^b$  induce on  $U(w_1(1))$  a left  $u(\mathfrak{m})^b$ -module structure via

$$\begin{aligned} e_1 \cdot u_1 &= u_1, & e_2 \cdot u_1 &= 0, & ae_1 \cdot u_1 &= u_2, & b^3 e_1 \cdot eu_1 &= 0, \\ e_1 \cdot u_2 &= 0, & e_2 \cdot u_2 &= u_2, & ae_1 \cdot u_2 &= 0, & b^3 e_1 \cdot u_2 &= 0, \\ e_1 \cdot u_3 &= u_3, & e_2 \cdot u_3 &= 0, & ae_1 \cdot u_3 &= 0, & b^3 e_1 \cdot u_3 &= u_2. \end{aligned}$$

The other elements of the basis of  $u(\mathfrak{m})^b$  act trivially.

## The functors

$$\begin{aligned} \mathrm{Ind}_e &:= {}_{\mathfrak{u}(\mathfrak{m})^b} \mathcal{M} \rightarrow {}_{\mathfrak{u}(\mathfrak{m})} \mathcal{M}, & \mathrm{Ind}_e(N) &= \mathfrak{u}(\mathfrak{m})e \otimes_{{}_{\mathfrak{u}(\mathfrak{m})^b}} N, \\ \mathrm{Res}_e &:= {}_{\mathfrak{u}(\mathfrak{m})} \mathcal{M} \rightarrow {}_{\mathfrak{u}(\mathfrak{m})^b} \mathcal{M}, & \mathrm{Res}_e(M) &= eM \end{aligned}$$

are inverse equivalences of categories.

## The functors

$$\begin{aligned}\mathrm{Ind}_e &:= {}_{\mathfrak{u}(\mathfrak{m})^b} \mathcal{M} \rightarrow {}_{\mathfrak{u}(\mathfrak{m})} \mathcal{M}, & \mathrm{Ind}_e(N) &= \mathfrak{u}(\mathfrak{m})e \otimes_{{}_{\mathfrak{u}(\mathfrak{m})^b}} N, \\ \mathrm{Res}_e &:= {}_{\mathfrak{u}(\mathfrak{m})} \mathcal{M} \rightarrow {}_{\mathfrak{u}(\mathfrak{m})^b} \mathcal{M}, & \mathrm{Res}_e(M) &= eM\end{aligned}$$

are inverse equivalences of categories.

Thus  $\mathrm{Ind}_e(U(w_1(1)))$  is the following 5-dimensional left  $\mathfrak{u}(\mathfrak{m})$ -module

$$e_1 \otimes u_1 \quad \xrightarrow{\quad} \quad e_2 \otimes u_2 \quad \xrightarrow{\quad} \quad ae_2 \otimes u_2 \quad \xrightarrow{\quad} \quad a^2e_2 \otimes u_2 \quad \xrightarrow{\quad} \quad e_1 \otimes u_3,$$

$\xleftarrow{\quad} \quad \xleftarrow{\quad} \quad \xleftarrow{\quad} \quad \xleftarrow{\quad}$

where the arrows oriented from left to right indicate the action of  $a$  while the arrows from right to left are the action of  $b$ .



## Category of finite-dimensional indecomposable left $u(\mathfrak{m})$ -modules

The non-isomorphic finite-dimensional indecomposable left  $u(\mathfrak{m})$ -modules are:

↪ the 8 families of string modules,

## Category of finite-dimensional indecomposable left $u(\mathfrak{m})$ -modules

The non-isomorphic finite-dimensional indecomposable left  $u(\mathfrak{m})$ -modules are:

- ↪ the 8 families of string modules,
- ↪ the 2 families of band modules,

## Category of finite-dimensional indecomposable left $u(\mathfrak{m})$ -modules

The non-isomorphic finite-dimensional indecomposable left  $u(\mathfrak{m})$ -modules are:

- ↪ the 8 families of string modules,
- ↪ the 2 families of band modules,
- ↪ the 2 simple modules and its respective projective covers.

Category of finite-dimensional indecomposable left  $u(\mathfrak{m})$ -modules

The non-isomorphic finite-dimensional indecomposable left  $u(\mathfrak{m})$ -modules are:

- ↪ the 8 families of string modules,
- ↪ the 2 families of band modules,
- ↪ the 2 simple modules and its respective projective covers.

## Next step

Determine the fusion rules. Precisely, for all finite-dimensional indecomposable left  $u(\mathfrak{m})$ -modules  $U$  and  $V$ , calculate the decomposition of  $U \otimes_{\mathbb{K}} V$  in direct sum of indecomposable modules.

The Drinfeld double  $D(H)$   
○○○○○

$u(\mathfrak{m})$  is tame type representation  
○○○○○

Indecomposable modules of  $u(\mathfrak{m})$   
○○○○○○●○

# References

## References

- [1] N. Andruskiewitsch, I. Angiono and I. Heckenberger. *On finite GK-dimensional Nichols algebras over abelian groups*. Mem. Amer. Math. Soc. **271**, No. 1329 (2021).
- [2] I. Assem, D. Simson, A. Skowroński. *Elements of the Representation Theory of Associative Algebras*. Lond. Math. Soc. Stud. Texts **65**, Cambridge Univ. Press (2006).
- [3] C. Cibils, A. Lauve, S. Witherspoon. *Hopf quivers and Nichols algebras in positive characteristic*. Proc. Amer. Math. Soc. **137** (12), 4029–4041 (2009) .
- [4] B. Wald and J. Waschbüsch *Tame biserial algebras*. J. Algebra **95**, 480-500 (1985).

The Drinfeld double  $D(H)$   
○○○○○

$u(\mathfrak{m})$  is tame type representation  
○○○○○

Indecomposable modules of  $u(\mathfrak{m})$   
○○○○○○○●

Thank you!